

1959

Variability in successive backcross generations of oats

Lester Norman Leininger
Iowa State University

Follow this and additional works at: <https://lib.dr.iastate.edu/rtd>



Part of the [Agriculture Commons](#), and the [Plant Sciences Commons](#)

Recommended Citation

Leininger, Lester Norman, "Variability in successive backcross generations of oats " (1959). *Retrospective Theses and Dissertations*. 2584.
<https://lib.dr.iastate.edu/rtd/2584>

This Dissertation is brought to you for free and open access by the Iowa State University Capstones, Theses and Dissertations at Iowa State University Digital Repository. It has been accepted for inclusion in Retrospective Theses and Dissertations by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

VARIABILITY IN SUCCESSIVE
BACKCROSS GENERATIONS OF OATS

by

Lester Norman Leininger

A Dissertation Submitted to the
Graduate Faculty in Partial Fulfillment of
The Requirements for the Degree of
DOCTOR OF PHILOSOPHY

Major Subject: Crop Breeding

Approved:

Signature was redacted for privacy.

In Charge of Major Work

Signature was redacted for privacy.

~~Professor-in-Charge, Farm Crops~~

Signature was redacted for privacy.

Head of Major Department

Signature was redacted for privacy.

Dean of Graduate College

Iowa State University
Of Science and Technology
Ames, Iowa

1959

TABLE OF CONTENTS

	Page
INTRODUCTION	1
REVIEW OF LITERATURE	2
MATERIALS AND METHODS	6
RESULTS AND DISCUSSION	11
Means and Variances	11
Selection	31
Linkage Study	36
GENERAL DISCUSSION	39
SUMMARY	41
LITERATURE CITED	42
ACKNOWLEDGMENT	45

INTRODUCTION

Backcrossing is a breeding system whereby the hybrid is crossed to one of the parental genotypes for one or more progeny generations. In general, this technique has two unique uses: (a) to add one or several simply inherited characters to an otherwise satisfactory strain, and (b) in testcrosses to determine genetic ratios, linkage and crossover percentages.

In a practical backcrossing program certain offspring which possess the desired characters are selected for further crossing to the recurrent parent. On the other hand, the theory of backcrossing is based upon unselected populations of infinite size.

It was the purpose of this study to determine whether the means and variances of an unselected^a sample of lines from several backcross generations would conform to those postulated by theory.

^aUnselected is used here to mean that no selection was knowingly practiced.

REVIEW OF LITERATURE

Backcrossing dates back to the time of the discovery of the particulate nature of inheritance by Mendel (18). Griffin as cited by Ware (36, p.667) used backcrossing in cotton breeding as early as 1867. Jennings (13) gave the quantitative results of the backcrossing system as follows:

If the original parents are AA and aa and the progeny are always mated back to one parent, the frequency of the recurrent parent genotype is $\frac{2^n - 1}{2^n}$, of the non-recurrent genotype is 0, and of the heterozygote is $\frac{1}{2^n}$ where n = number of backcross generations. Briggs (2), and Riddle and Baker (25) later expanded the formula of Jennings (13) to include the number of factor pairs as follows:

$$\% \text{ homozygosity} = \left(\frac{2^m - 1}{2^m} \right)^n \quad \text{where: } m = \text{number of backcross generations and}$$


n = number of heterozygous factor pairs.

Wright (37) using the method of path coefficients arrived at an identical solution.

It remained, however, for Harlan and Pope (9) to point out the place of backcrossing in cereal breeding with experiments involving the smooth awn character in barley. This work stimulated the thought of many plant breeders. Richey (23) published the theory of convergent improvement, which is essentially reciprocal backcrossing, in corn. Later, Richey and Sprague (24), Murphy (19), and Hayes et al. (12) reported that convergent improvement was successful in improving inbred lines of corn.

Mather (17, pp. 68-71) published the biometrical expectations of

reciprocal backcrosses. Using the following model:



He states "the contributions of gene $A - a$ to the mean of the two backcrosses, as measured from the midparent, will be $B_1 = 1/2(d_a + h_a)$ and $B_2 = 1/2(h_a - d_a)$, where B_1 is the backcross to the AA parent which gives the plus deviation, and B_2 that to the aa parent. The contributions of $A - a$ to the variances of the backcrosses are $B_1 = 1/4(d_a - h_a)^2$ and $B_2 = 1/4(d_a + h_a)^2$. The contributions of d and h to these variances cannot be separated as they stand, but if the variances are summed, $A - a$ contributes $1/2(d_a^2 + h_a^2)$ to the total, and the contributions of d and h can then be separated."

The primary advantage of backcrossing over selfing is that a higher percentage of homozygotes result (2). In the F_2 of a single cross, either parent is expected once in 4^n plants, whereas in the Bc_1 , the recurrent parent is expected once in 2^n number of plants. To quote Knight (15, p. 77) "the essential value of backcrossing is that it provides a mean of limiting the heterogeneity which would result from 'straight' crosses between two types, making it possible to produce a hybrid similar to whichever of the two varieties has the more valuable genetic constitution, yet containing desirable characters transferred from the other parent. Backcrossing obviates the necessity for rigid selection generation after generation in F_2 , F_3 , F_4 , etc., by progressively and

automatically rendering the hybrid more and more homogeneous."

Briggs (1) reported success in transferring bunt resistance from the Martin to Baart variety of wheat. He (3) reported excellent progress in producing better wheat varieties with the backcross system, and proposed the use of backcrosses to form isogenic lines to determine the effect of different genes on yield.

Briggs and Allard (4) state three basic requirements for a backcross breeding program: (a) a satisfactory recurrent parent, (b) retention of the intensity of the character under transfer through several backcrosses, and (c) the genotype of the recurrent parent must be reconstituted in a reasonable number of backcrosses with small populations. The expression of a gene may vary with different genetic backgrounds according to Simpson and Weindling (28) and Smith (29) and apparently there is no way of predicting the gene expression on a given background without trying it.

Others who have successfully used the backcross system on agricultural crops are Emsweller and Jones (6), Johnson and Hayes (14), Suneson et al. (31), Suneson (30) and Valteau (34,35). Thomas (33) has given a comprehensive review of backcrossing in tobacco, tomato, cotton and potatoes.

Backcrossing appears to have little use in transferring quantitative characters unless they are controlled by only one or two major genes. However, the California workers were successful in transferring genes for maturity and height from Romona to Baart wheat, and Powers (21,22) was able to transfer quantitative characters in tomatoes. Everson (?) and Suneson et al. (32) reported a major yield gene in barley which could

be transferred by backcrossing since it was linked to genes for rachilla hair length and awn barbing.

Mather (16) calculated for several probability levels the number of backcross progeny necessary to assure the breeder that the gene being transferred will be present in the population. The use of these tables was illustrated by Harrington (10) and Hayes et al. (11).

The usefulness of the backcross technique in studying linkage relationships of genes and chromosome mapping has been demonstrated by Burnham and Cartledge (5). According to Briggs and Allard (4, p. 134) the backcross method is more effective than selfing in breaking linkages between desirable and undesirable genes. They state "in any well executed backcross program, undesirable linkages may be expected to be broken in a portion of the derived lines, even if the crossover value is as small as 1 percent and the undesirable allele is not expressed."

MATERIALS AND METHODS

The recurrent and non-recurrent oat varieties used in this study were Cherokee and P.I. 185785, respectively. Cherokee (C.I. 5444) was selected from the cross D69 x Bond. It is early maturing, short in height, susceptible to race 202 of crown rust, and produces plump, white grain of good test weight. P.I. 185785 (C.I. 6952) was introduced into North America from Uruguay. It is late maturing, tall in height, resistant to race 202 of crown rust and produces long, slender, red grain of low test weight.

Cherokee and P.I. 185785 are both allo hexaploid varieties with 21 pairs of chromosomes (20,27). Further, both parents are self pollinated and are considered to be homogeneous lines.

The F_1 between Cherokee and P.I. 185785 was made in 1953 resulting in 13 hybrid seeds. Plants from these F_1 seeds were backcrossed to Cherokee to give the Bc_1^a generation which in turn was backcrossed to Cherokee, etc., until 4 backcross generations were available. In addition to backcrossing to each plant, F_2 seed was also saved and stored so that all crosses could be grown in a common yield test. The number of lines in each backcross generation is shown in table 1.

Pedigrees were kept so that each oat strain could be traced to the proper Bc_1 , Bc_2 , Bc_3 or Bc_4 family. This was necessary so that the

aBc_1 = P.I. 185785 x Cherokee² = first backcross
 Bc_2 = P.I. 185785 x Cherokee³ = second backcross
 Bc_3 = P.I. 185785 x Cherokee⁴ = third backcross
 Bc_4 = P.I. 185785 x Cherokee⁵ = fourth backcross.

Table 1. Number of progenies tested in each backcross generation.

Backcross no.	No. of progenies
1st	25
2nd	110
3rd	215
4th	275

variances could be assigned to the proper within and between family categories. The pedigrees of all lines are shown in figure 1.

The crossing was completed in 1956 and the seed for the 4 backcross generations grown for increase at Aberdeen, Idaho in 1957. An experiment including 625 backcross lines (in the F_2 or F_3 generation), P.I. 185785 entered 10 times, and Cherokee 17 times, making a total of 652 entries, was grown in 1958. The experimental design was a randomized block with 8 replications, 4 grown at Ames and 4 at Kanawha, Iowa. The plot size was one hill with a 1-foot spacing between hills (26). Thirty seeds were sown in each plot which was equivalent to a planting rate of 3 bushels per acre.

Each replication was 30 x 22 feet including 8 dummy plots to bring the number to 660 per replication. At each location, the experiment was surrounded by 3 rows of hills to serve as competition and border.

Growing conditions for oats were good to excellent at Ames and Kanawha in 1958. The rainfall distribution was variable with a short supply early in the growing season and an abundant supply in the later

Figure 1. The complete hierarchy of all families through four successive backcross generations.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
84

1
2
3
4
5

1
2
3
4
5
6

[illegible]

[illegible]

part. The plots at Ames received one sprinkler irrigation early in the season. Since the experiment included lines both resistant and susceptible to crown rust and other foliar diseases, it was sprayed with a fungicide (Zineb) so that disease reaction would not confound the expression of the characters measured.

The measurements taken were date of heading and plant height on the 4 replications grown at Ames, and grain yield on all 8 replications. Weight-per-volume was measured as 2 replications since it was necessary to combine the seed of an entry from four replications to order to have enough to take a weight-per-volume measurement. These measurements were taken in a glass cylinder measuring 7.7 mm in diameter and 3.9 in depth. Date of heading was recorded when 50 percent of the heads in a plot were completely emerged from the boot; plant height was measured in inches from ground level to the panicle tips; and grain yield was recorded in grams per plot. Each of these characters is multigenically inherited and subject to continuous distribution, both characteristic of quantitative characters.

RESULTS AND DISCUSSION

Means and Variances

The mean squares from the analysis of variance of the data for grain yield, date of heading, plant height and weight per volume are presented in table 2.

All of the mean squares involving lines within generations, except for the parental entries, were highly significant. While the general trend was for the mean squares to be reduced with a greater number of backcrosses, the reduction was not pronounced until the Bc_4 . The mean squares for height and weight-per-volume were quite constant in the Bc_1 , Bc_2 and Bc_3 generations but reduced in the Bc_4 . For some unknown reason, yield and date of heading mean squares were low in the Bc_2 and Bc_4 but high in the Bc_3 generation. One or two Bc_1 families caused most of this abnormality.

The mean squares for the two parents approximated the error variance. P.I. 185785 showed significant variation for date of heading but this may have been due to its poor adaptation to Iowa growing conditions.

One of the important factors in a backcrossing program is the rapidity with which the performance of the hybrid lines return to the level of the recurrent parent. This can be studied in two ways: (a) by the means and variances of the different generations and (b) by the range of performance among lines within backcross generations.

The mean squares for Cherokee vs. the means of the backcross generations are presented in table 3. Since the Cherokee parent contributes one-half of the germplasm to each backcross, the mean squares measuring

Table 2. Mean squares for four quantitative characters among lines within generations.

Source of variation	Degrees of freedom	Characters			
		Yield	Date	Height	Wt./vol.
Entries	649	165**	99**	13**	21**
Entries x loc.	649	37	--	--	--
Between generations	5	2133**	7558**	468**	486**
Among entries in generation	644	150**	41**	10**	18**
Among Bc ₁ entries	24	277**	138**	12**	26**
Among Bc ₂ entries	108	127**	33**	12**	22**
Among Bc ₃ entries	213	198**	80**	11**	22**
Among Bc ₄ entries	274	121**	9**	8**	12**
Among Cherokee entries	16	27	0	2	2
Among P.I. 185785 entries	9	28	3*	1	--
Error	--- ^a	26	1.4	2	5

^aD.f. for error are as follows: Yield = 3894; Date = 1953; Height = 1953; and Wt./vol. = 637.

For this and all following tables:

*Exceeds the 5% level of probability

**Exceeds the 1% level of probability

differences between Cherokee and the backcross means should decrease with successive generations of backcrossing. All mean squares were significant, and date of heading, height and weight-per-volume mean squares decreased as expected, especially in the Bc₄. However, the trend for yield was for the mean squares to increase.

The mean performance for the parental strains (table 4) differed widely for each of the characters measured. In the backcross material, the means for date of heading, height and weight-per-volume progressed toward the Cherokee parent gradually (figure 2, table 4), but perhaps not as rapidly as expected since by Bc₄ the germplasm contribution from

Table 3. Mean squares for Cherokee vs. the backcross generation means for four quantitative characters.^a

Source of variation	Degrees of freedom	Characters			
		Yield	Date	Height	Wt./vol.
Between generations	5	2133**	7558**	468**	486**
Cherokee vs. Bc ₁	1	518**	554**	350**	539**
Cherokee vs. Bc ₂	1	1102**	95**	466**	697**
Cherokee vs. Bc ₃	1	1015**	67**	297**	592**
Cherokee vs. Bc ₄	1	1357**	28**	259**	182**
Cherokee vs. P.I. 185785	1	3100**	25,301**	1840**	---

^aNon orthogonal breakdown of sum of squares.





Table 4. Means of four quantitative characters for the parents and four backcross generations.

Parents and backcross generations	No. of entries	Means			
		Yield	Date	Height	Wt./vol.
Entries	652	32	7.3	35	70
Bc ₁	25	32	8.8	36	68
Bc ₂	110	32	7.0	36	69
Bc ₃	215	32	6.8	35	69
Bc ₄	275	32	6.5	35	71
Cherokee	17	29	5.8	33	74
P.I. 185785	10	21	37.5	41	62* ^a

^aThis is an estimate over all eight replications due to a short supply of seed.

Cherokee to the backcross plants should be nearly 97 percent. The theoretical 3 percent contributed by the P.I. 185785 parent was apparently very potent in expression or it is possible that some unknown selection pressure was operating which may have caused retention of a larger percentage of the P.I. 185785 germplasm than was expected.

Figure 2. Actual and theoretical means for four quantitative characters in the backcross generations.







Also included in figure 2 are the expected means for the four quantitative characters in each backcross generation. The theoretical curve was constructed on the assumption of additive gene action with the midparent value representing the F_1 and the Cherokee value as the performance after an infinite number of generations of backcrossing. The expected progress toward the Cherokee value was one-half of the remainder for each succeeding generation. The trend in mean values for the weight-per-volume suggest dominance for low weight-per-volume contributed by the P.I. 185785 parent. In contrast, the actual curve of means for heading date suggest dominance of earliness from the Cherokee parent.

The mean yield per plot was approximately 32 grams for all four backcross generations. This value was significantly above both parental strains. Such a heterotic effect might be expected in the early backcross generations since yield is subject to geometric epistasis (8). However, the yield nevertheless should have regressed toward the recurrent parent mean to a certain extent with four backcrosses. Since yield is probably affected by a large number of genes, the proportion of individuals which would be reconstituted like Cherokee would be small even in the Bc_4 . This suggests that there would be ample opportunity for considerable geometric epistasis as late as the Bc_4 and, therefore, the yield level of the backcross generations would not regress very rapidly. The curve of height means is somewhat parallel to the theoretical curve but at a lower level.

The theoretical and actual total variances for the four characters in each generation are plotted in figure 3. The theoretical curves for

Figure 3. Actual and theoretical variances for four quantitative characters in four backcross generations.



yield and date of heading were constructed on the basis of epistatic effects, whereas those for height and weight-per-volume were based on dominance or additive gene action (the reason to be discussed in conjunction with tables 10 and 11). For construction of the theoretical curves, the true variance was assumed to be the same as that of the Bc_1 . Needless to say, this assumption may be subject to considerable error, but it should not affect comparisons where trends of actual and theoretical curves are involved.

The theoretical curves for epistasis were calculated on two and three gene models with varying gene effects. The expected reduction of total variance for succeeding backcross generations was less than one-half and the variance reductions from one generation to the next were less with epistasis than with dominance or additive gene action. In all four characters the total variances were reduced with continued backcrossing.

For height and weight-per-volume, the actual variances were not reduced as rapidly as expected. The variances for yield and date of heading were sporadic. As noted earlier, the Bc_3 variances were larger than those in Bc_2 . The only reasonable explanation for the large Bc_3 variance is that a divergent sample of gametes was obtained when the backcross was made on the Bc_2 .

Tables 5, 6 and 7 contain the mean squares within individual Bc_1 families for the Bc_2 , Bc_3 and Bc_4 generations. While most Bc_4 families are less variable than their Bc_3 progenitors, several Bc_3 families were more variable than their Bc_2 ancestors.

Part of the explanation for variances higher than expected in Bc_3

Table 5. Mean squares for four quantitative characters within Bc₁ families in the Bc₂ generation.

Source of variation	Degrees of freedom	Characters			
		Yield	Date	Height	Wt./vol.
Bc ₂ generation	109	127**	33**	12**	22**
Within family <u>a</u>	3	157**	240**	12**	20**
Within family <u>b</u>	--	--	--	--	--
Within family <u>c</u>	1	30	3	12**	36**
Within family <u>d</u>	4	87	6**	7**	9
Within family <u>e</u>	1	12	0	6	0
Within family <u>f</u>	1	663**	1	12**	0
Within family <u>g</u>	9	235**	33**	13**	26**
Within family <u>h</u>	4	13	3	2	4
Within family <u>i</u>	1	53	2	2	1
Within family <u>j</u>	4	75*	1	6*	21**
Within family <u>k</u>	12	125**	23**	13**	16**
Within family <u>l</u>	3	123**	12**	14**	7
Within family <u>m</u>	2	31	26**	21**	29**
Within family <u>n</u>	3	46	1	0	29**
Within family <u>o</u>	2	92**	18**	1	4
Within family <u>p</u>	4	12	1	2	25**
Within family <u>q</u>	3	76*	2	7*	13
Within family <u>r</u>	3	28	25**	9**	46**
Within family <u>s</u>	3	13	4*	1	14*
Within family <u>t</u>	1	1	10**	18**	6
Within family <u>u</u>	4	174**	14**	16**	15*
Within family <u>v</u>	7	30	1	2	12*
Within family <u>w</u>	8	53*	2	5**	13*
Within family <u>z</u>	4	150**	3	3	9
Between Bc ₂ families	22	238**	89**	30**	41**

families may be due to the fact that P.I. 185785 was not adapted to Iowa growing conditions, and the measurements taken directly on this variety did not express its true potential of performance. When a portion of the P.I. 185785 germplasm was substituted onto an essentially Cherokee background, it had an opportunity for a maximum expression. Another contributing factor may be a divergent "gametic sample" in the Bc₂.

Table 6. Mean squares for four quantitative characters within Bc_1 families in the Bc_3 generation.

Source of variation	Degrees of freedom	Characters			
		Yield	Date	Height	Wt./vol.
Bc_3 generation	214	198**	80**	11**	22**
Within family <u>a</u>	6	125**	167**	33**	42**
Within family <u>b</u>	6	31	9**	9**	5
Within family <u>c</u>	13	118**	14**	2	16**
Within family <u>d</u>	19	164**	6**	4**	26**
Within family <u>e</u>	2	683**	3	9*	43**
Within family <u>f</u>	5	290**	108**	15**	23**
Within family <u>g</u>	8	1121**	1018**	31**	36**
Within family <u>h</u>	15	236**	13**	5**	15**
Within family <u>i</u>	7	113**	3	3	8
Within family <u>j</u>	9	105**	2	7**	18**
Within family <u>k</u>	30	86**	25**	12**	13**
Within family <u>l</u>	10	336**	44**	28**	23**
Within family <u>m</u>	14	146**	2	6**	13**
Within family <u>n</u>	1	81	3	3	2
Within family <u>o</u>	5	22	1	2	1
Within family <u>p</u>	3	43	2	1	2
Within family <u>q</u>	6	67*	3*	6**	5
Within family <u>r</u>	1	0	2	0	30**
Within family <u>s</u>	16	48*	1	2	9*
Within family <u>t</u>	2	2	1	1	11
Within family <u>u</u>	--	--	--	--	--
Within family <u>v</u>	5	73*	3	2	2
Within family <u>w</u>	1	1	0	1	25**
Within family <u>z</u>	8	76**	2	2	6
Between Bc_3 families	22	422**	252**	33**	78**

To illustrate, the major source of Bc_2 variation in family a (table 8 and figure 4) was due to entry 027 whose mean heading date was from two to four times larger than the other Bc_2 entries. (The numbers enclosed parenthetically are entry numbers and the adjacent numbers are the dates of heading.) This shows the gametic variability available from the Bc_1 . Entry 027 gave one gamete which produced a late zygote

Table 7. Mean squares for four quantitative characters within Bc_1 families in the Bc_4 generation.

Source of variation	Degrees of freedom	Characters			
		Yield	Date	Height	Wt./vol.
Bc_4 generation	274	121**	9**	8**	12**
Within family <u>a</u>	12	95**	20**	15**	4
Within family <u>b</u>	6	55*	5**	4*	4
Within family <u>c</u>	11	45	1	3	6
Within family <u>d</u>	3	47	4*	1	2
Within family <u>e</u>	5	313**	1	3	1
Within family <u>f</u>	2	2	1	2	2
Within family <u>g</u>	11	103**	25**	24**	33**
Within family <u>h</u>	7	108**	6**	2	8
Within family <u>i</u>	8	74**	3	4*	4
Within family <u>j</u>	63	91**	2	4**	7
Within family <u>k</u>	55	28	2	7**	5
Within family <u>l</u>	17	120**	7**	11**	5
Within family <u>m</u>	9	157**	1	4*	14**
Within family <u>n</u>	1	14	6*	3	9
Within family <u>o</u>	2	53	2	6*	4
Within family <u>p</u>	0	0	0	0	0
Within family <u>q</u>	1	6	0	0	2
Within family <u>r</u>	5	47	0	5*	6
Within family <u>s</u>	13	199**	4**	5*	12**
Within family <u>t</u>	5	62*	0	3	5
Within family <u>u</u>	--	--	--	--	--
Within family <u>v</u>	2	50	3	1	1
Within family <u>w</u>	1	1	0	6*	1
Within family <u>z</u>	13	32	2	3*	11*
Between Bc_4 families	22	570**	638**	33**	62**

and one which was early, so the genetic range was still highly variable for date of heading genes in the Bc_2 . Although the Bc_2 should have contained seven-eighths Cherokee genotype, segregation produced gametes which were quite unlike Cherokee. However, it is evident that the sample of gametes obtained in the Bc_3 generation contained an abnormally high number of "non-Cherokee" gametes. In the Bc_3 , the proportion of the "rare" gametes should have become small enough so that the sample

Table 8. Mean squares for date of heading for lines within family a.


Source of variation	Degrees of freedom	Characters			
		Yield	Date	Height	Wt./vol.
Bc ₂					
Family <u>a</u>	3	157**	240**	12**	20**
Bc ₃					
Family <u>a</u>	6	125**	167**	33**	42**
Within Bc ₃ family 1	1	5	13**	3	4
Within Bc ₃ family 2	1	324**	406**	41**	21**
Within Bc ₃ family 3	1	68	0	2	1
Between Bc ₃ families	3	118**	194**	50**	76**
Bc ₄					
Family <u>a</u>	12	95**	20**	15**	4
Within Bc ₄ family 1	2	30	1	3	2
Within Bc ₄ family 2	9	73**	2	7**	3
Between Bc ₄ families	1	425**	217**	118**	17

included in this study would not be expected to include them very often; and they were only occasionally found in the Bc₄ progeny. This was more true for date of heading and yield than for the other two characters.

Since backcross theory is based only upon the assumption of additive genetic effects, it is further possible that genetic systems such as dominance, epistasis, linkage, etc. could have produced the observed variability within families of the second and third backcross generations. However, just what effect such genetic systems would have on backcross populations has not been established.

The mean squares between Bc₁ families in the Bc₂, Bc₃ and Bc₄ for

Figure 4. Entry numbers (in parentheses) and means for dates of heading for progenies in family a.



the four quantitative characters are given in table 9. The variation between families did not change significantly from one generation to the next for height and weight-per-volume, but it did increase significantly for the other two characters, date of heading and grain yield. The magnitude of increase of the variances between Bc_1 families was rapid for these latter two characters. The F test showed that the Bc_4 variance for yield was significantly greater than the Bc_2 variance. For date of heading, the variance between families was significantly greater in each successive backcross.

The trend expected for between family variances over generations was determined using a two gene model based upon different types of gene action (table 10). The non-recurrent parent was represented by aaBB and the recurrent parent by AAbb, and the genes were given the following effects: $A = 3$; $B = 5$; $a = b = 1$. The variances between Bc_1 families should have remained constant over all backcross generations with either dominance or additive genetic effects. Since the variances between Bc_1 families for height and weight-per-volume did not differ significantly, the genes affecting these two characters probably displayed dominance and/or interacted additively. The variances between Bc_1 families for yield and date of heading increased with successive backcross generations indicating epistatic gene action. For yield, this was expected since it was subject to geometric epistasis, but previous studies on date of heading have given no indication of epistasis.

Transgressive segregation is frequently observed for quantitative character in the progeny of crosses in cereals, but generally not when the parents are as diverse as those used in this study. Such observations

Table 9. Mean squares between Bc_1 families for quantitative characters in the Bc_2 , Bc_3 and Bc_4 .

Source of variation	Degrees of freedom	Characters			
		Yield	Date	Height	Wt./vol.
Between Bc_2 families	22	238**	89**	30**	41**
Between Bc_3 families	22	422**	252**	33**	78**
Between Bc_4 families	22	570**	638**	33**	62**

Table 10. Theoretical variances between Bc_1 families in four backcross generations with epistasis, dominance or additive gene action, using a two gene model and selfing to form F_2 .

Generation	Variances between Bc_1 families ^a		
	Epistasis	Dominance	Additive
Bc_1	23.40	3.08	6.67
Bc_2	24.25	3.08	6.67
Bc_3	25.56	3.08	6.67
Bc_4	26.26	3.08	6.67

^aThe effect of the genes are as follows: $A = 3$; $B = 5$; $a = b = 1$.

are expected even less in backcross progeny. Be this as it may, transgressive segregation was observed as late as the Bc_3 . Date of heading data illustrates this phenomena, although it is exhibited by the other characters also.

In the Bc_3 generation of family g (figure 5), some entries transgressed the parental range for date of heading. Table 11 contains the

Figure 5. Entry numbers (in parentheses) and dates of heading for progenies in family g.




Table 11. Mean squares for date of heading for lines within family g.

Source of variation	Degrees of freedom	Characters			
		Yield	Date	Height	Wt./vol.
Bc ₂					
Family <u>g</u>	9	235**	33**	13**	26**
Bc ₃					
Family <u>g</u>	8	1121**	1018**	31**	36**
Within Bc ₃ family 1	3	38	1	1	20*
Within Bc ₃ family 2	2	1393**	1211**	7*	--
Between Bc ₃ families	3	2023**	1905**	115**	52**
Bc ₄					
Family <u>g</u>	11	103**	25**	15**	33**
Within Bc ₄ family 1	3	25	2	2	11
Within Bc ₄ family 2	6	6	18**	24**	5
Between Bc ₄ families	2	512**	81**	8*	150**

mean square analysis for these data. The heading date of entry 241 was 43.3 compared to 37.5 for P.I. 185785. In contrast, entry 243 was earlier than Cherokee (5.8). The extremely late lines were unexpected since the gametes which contributed to the late zygotes in the Bc₃ were produced from a fairly early Bc₂ plant. Aberrant gametic sampling seems to be the only possible explanation for this phenomenon. The Bc₃ entries 106, 241 and 242 with heading dates of 42.5, 43.3 and 12.8, respectively, were progenies from the cross Cherokee x Bc₂ entry 041 for which the heading dates were 5.8 and 10.3, respectively. While errors of pedigree and classification are possible in any experiment, this does not appear to be the case here since entries 106 and 241 both trace to the same parents.

Selection

From a practical standpoint, the rapidity with which the mean values for various characters in the backcross material returned to the Cherokee parental values was of limited importance. Even though the Bc_4 mean values for yield, height and weight-per-volume were considerably divergent from Cherokee, some lines should have been nearly identical to the recurrent parent. Selection for the recurrent parent type would be easier with additional backcrosses.

In the field, it would be possible to select plants for continued backcrossing which were more like the recurrent parent than the average of the hybrid population. For example, plants could have been selected which were most like Cherokee for date of heading and height before it was necessary to cross. Yield and weight-per-volume could be determined only when the plants were ripe; so, selection for these two characters would result in the loss of a year's time in crossing. Therefore, it would be most expedient to select for the first two characters only.

Five families were selected in the first backcross which were nearly the same as Cherokee for height and heading date. These families were compared with Cherokee through the four backcross generations (table 12). Selection for Bc_1 lines which were like Cherokee for heading date and height was effective in retaining these two characters near the Cherokee phenotype in the three subsequent backcross generations. The mean yield of the selected lines was closer to Cherokee than in the unselected families; but the reverse was true for weight-per-volume. The mean squares for "Cherokee vs. selected families" were considerably

Table 12. Mean squares from analysis of Cherokee vs. selected families and Cherokee vs. unselected families in the Bc₂, Bc₃ and Bc₄.

Source of variation	Degrees of freedom	Characters			
		Yield	Date	Height	Wt./vol.
Bc ₂					
Between selected families Cherokee vs. selected families	3	171*	14	1	30
Cherokee vs. unselected families	1	41	1	96	309**
Among lines in selected families	1	1556**	117**	514**	703**
Among lines in unselected families	12	58	10	10	18
	75	106	21	8	18
Bc ₃					
Between selected families Cherokee vs. selected families	3	306	7	14	73
Cherokee vs. unselected families	1	258	1	134**	1037**
Among lines in selected families	1	1129**	85	311**	465**
Among lines in unselected families	25	186	3	6	12
	167	170	69	9	18
Bc ₄					
Between selected families Cherokee vs. selected families	3	365	4	1	21
Cherokee vs. unselected families	1	41	2	56**	355**
Among lines in selected families	1	1558**	31**	277**	156**
Among lines in unselected families	20	220	3	1	11
	232	70	5	7	7

smaller than those for "Cherokee vs. unselected families" in both of the selected characters and yield. The most likely explanation for the selection of Cherokee type backcross lines for heading date and height being effective in obtaining types which are also like Cherokee in yield would be linkage drag. Genes which were responsible for the "Cherokee heading date and height" were linked with genes for yield and therefore, selection for the first two characters "dragged" along many Cherokee yield genes.

When oat plants were grown in the greenhouse, height was not very heritable, so this character could not be used to select Cherokee type plants with good precision. However, date of heading is heritable in the greenhouse and could be used as an effective selection criterion. Since most of the oat crossing is done in the greenhouse, the above idea was pursued in a selection experiment. Three Bc_1 families were selected, one early, one like Cherokee, and one late for date of heading. The means and ranges for these families in each generation are presented in table 13. In many cases, the family range for a character was smaller than the recurrent parent range, especially in the Bc_4 . In general, selection for either early (family c) or late (family f) heading date was detrimental to mean yield although family c exceeded the recurrent parent in yield in the Bc_4 . Height regressed toward the recurrent parent quite rapidly.

Selection for the recurrent parent heading date in family f after the Bc_1 would have missed the entry in the Bc_2 which yielded 43 grams per plot. This entry produced progeny yielding up to 39 grams in the Bc_3 but was not used to produce Bc_4 progeny. Selection for the recurrent

Table 13. Mean and range for four quantitative characters in each backcross generation for three families with selection for heading date in each backcross.

Backcross generation	Mean				Range			
	Yield	Date	Height	Wt./vol.	Yield	Date	Height	Wt./vol.
Family c								
Bc ₁	28	2.0	35	72	---	----	---	---
Bc ₂	28	4.2	34	68	26-29	3.5-4.8	33-36	65-71
Bc ₃	27	5.9	34	68	25-29	5.3-6.5	33-34	67-68
Bc ₄	33	5.9	34	69	31-38	5.5-6.8	33-35	68-71
Family f								
Bc ₁	23	23.8	39	65	---	----	---	---
Bc ₂	36	11.8	38	65	30-43	11.3-12.0	37-40	65-66
Bc ₃	24	6.5	35	67	---	----	---	---
Bc ₄	20	7.3	34	68	19-20	7.0-7.5	34-35	67-69
Family g								
Bc ₁	37	5.3	36	72	---	----	---	---
Bc ₂	33	5.5	34	71	31-34	4.5-6.8	34-35	68-74
Bc ₃	33	6.1	34	73	31-34	5.5-6.8	34-35	71-75
Bc ₄	31	6.3	33	73	30-32	5.8-6.8	33	72-73
Cherokee								
	29	5.8	33	74	26-33	5.5-6.3	32-34	72-75

parent heading date in family g caused all attributes to regress the recurrent parent phenotype.

In general, it appears that selection for date of heading and height, if possible, would result in recapturing the recurrent parent characteristics more rapidly than if random selection of plants for backcrossing were practiced.

Another way to measure whether selection might be effective would be through the correlations between generations. The correlation coefficients of family means for all combinations of the four backcross generations for yield, height, date of heading and weight-per-volume are given in table 14. For date of heading, the correlations were significant and very high for all comparisons. This further indicates that selection for date of heading in one backcross generation would be effective in returning the backcross progenies to the Cherokee type more rapidly than if crosses were made at random. All of the height correlations were significant, but not as high as those for date of heading. However, selection for height in any or all of the backcross generations should make the regression toward the recurrent parent more rapid.

The correlations between Bc_2 , Bc_3 , and Bc_4 for weight-per-volume were highly significant indicating a good relationship between these generations for this character. However, the Bc_1 was significantly correlated with only the Bc_2 . Although several of the correlation coefficients for yield were significant the relationship between generations for this character were, in general, not very good.

Although with a sufficient number of backcross generations the plant breeder is assured of recovering the recurrent parent phenotype,

Table 14. Correlation coefficients among four backcross generations for four characters.

	Bc ₂	Bc ₃	Bc ₄	Bc ₂	Bc ₃	Bc ₄
	<u>Yield</u>			<u>Date</u>		
Bc ₁	.30	.57**	.45*	.87**	.87**	.69**
Bc ₂		.23	.13		.82**	.87**
Bc ₃			.55**			.72**
	<u>Height</u>			<u>Wt./vol.</u>		
Bc ₁	.76**	.52*	.57**	.65**	.37	.40
Bc ₂		.73**	.62**		.59**	.75**
Bc ₃			.48*			.61**

any procedure which would reduce the recovery time would be desirable. Selection of backcross lines in each generation which are most like the recurrent parent may reduce the number of backcrosses necessary to attain the desired goal. This study indicated that phenotypic expression of one backcross generation is well correlated with the next generation for date of heading and height. Furthermore, selection for the "Cherokee type" expression of these characters also resulted in the yield of the selected lines being closer to the Cherokee performance than was the case in the backcross population as a whole.

Linkage Study

Backcrossing has been used frequently in oats to transfer rust resistance into adapted varieties. In backcrossing to add rust resistance, selection for resistant progenies is essential in order to keep

the population size manageable while retaining progeny that carry a worthwhile intensity of resistance. If desirable or undesirable genes from the non-recurrent parent were closely linked to the resistance gene, selection for rust resistance would be either beneficial or harmful, respectively, to the primary objective. A beneficial example was found by Suneson et al. (32) where high yield genes were closely linked with the semi-smooth awn (*rr*) gene in barley.

Since the non-recurrent parent used in this study was so poorly adapted in Iowa, it seemed likely that only undesirable genes would be linked to the crown rust resistance gene being transferred to Cherokee. If such were the case, and if linkage were important, the mean performance for the four characters measured would not be as close to Cherokee in the resistant as in the susceptible backcross lines.

In the Bc_1 , 8 entries were resistant to crown rust race 202 and 15 were susceptible. In the Bc_2 and Bc_3 (table 15), the resistant and susceptible lines were not significantly different for any of the four agronomic characters measured. In the Bc_4 , the means of the resistant and susceptible groups were significantly different for date of heading, height, and weight-per-volume. This significance could be important since the means of the resistant lines were nearer to Cherokee than were the susceptible line means. However, the differences were very small, 0.6 of an inch in height, 0.3 of a day in heading date, 0.5 of a gram in weight-per-volume, and probably of little consequence. Apparently, detrimental or beneficial genes from the non-recurrent parent which might be linked to the gene for crown rust resistance were either non-existent or affected the four characters measured very little.

Table 15. Mean squares from analysis of resistant vs. susceptible families in the Bc₂, Bc₃ and Bc₄.

Source of variation	Degrees of freedom	Characters			
		Yield	Date	Height	Wt./vol.
Bc ₂					
Total	109	127*	33**	12*	22
Resistant vs. susceptible	1	30	33	3	79
Between resistant families	7	176	74**	25**	38*
Between susceptible families	14	285*	101**	35**	46**
Among lines in families	87	98	19	8	17
Bc ₃					
Total	214	198	80*	11*	22*
Resistant vs. susceptible	1	239	6	3	30
Between resistant families	7	344	251**	51**	94**
Between susceptible families	14	477**	241**	27**	72**
Among lines in families	192	172	62	9	16
Bc ₄					
Total	274	121**	9**	8*	12**
Resistant vs. susceptible	1	48	205**	85**	322**
Between resistant families	7	669**	18**	13*	52**
Between susceptible families	14	559**	77**	39**	56**
Among lines in families	252	82	4	6	8

GENERAL DISCUSSION

In this study, the means and variances of quantitative characters in oats did not return to the recurrent parent by a factor of one-half of the remaining difference between the backcross and recurrent parent means in each succeeding generation. However, the means and variances did, in general, return toward the recurrent parent, although at varying rates for the different characters. Any or all of the following factors may aid in explaining the observed discrepancies:

First, backcross theory is based upon infinite populations, and while the population size in each generation of this study was large compared to a practical backcrossing program, it was small in comparison to infinity. Therefore, it is possible that this "sample" was quite divergent from the infinite population.

Second, diploidy has been assumed in all backcross theory. While some inheritance in oats is diploid in nature, it is unlikely that genes responsible for quantitative characters act as a diploid completely since oats is an allohexaploid species.

Third, theory is based upon such assumptions as additive interaction of loci, complete dominance, equal frequency alleles, equal gene effects between loci, no linkage and no epistasis. Certain of these assumptions are not correct for oats. For example, this study shows that the different quantitative characters are probably controlled by different inter-loci reaction systems.

Fourth, the return to the recurrent parent is usually calculated as percent homozygosity (or heterozygosity) which leads to a reduction by

one-half per backcross generation. Assuming additive, the means return to the recurrent parent by one-half per generation but this is unlikely to be the only genetic system operative. Simple gene models indicate that the variance is reduced in early generations by a factor of less than one-half.

While the number of backcross generations used in this study was too limited to supply a conclusive solution to the use of backcrossing for quantitative characters, encouragement for the plant breeder is found in the results. The means of all characters measured regressed toward the recurrent parent, except yield which showed a persistent increase over the recurrent parent. If the increase for yield should continue through further backcrosses while other characters returned to the recurrent parent phenotype, backcrossing would be useful not only for transferring qualitative characters, but also for producing higher yielding varieties as well. The reduction in yield variance without a reduction in mean yield indicated this to be a possibility.

SUMMARY

In four generations of backcrossing in oats, the mean date of heading, plant height, weight-per-volume and grain yield regressed toward the recurrent parent, but at rates different than expected on the basis of additive gene action alone. Plant height and weight-per-volume returned at a slower rate than expected; date of heading returned faster than expected. Mean yield showed considerable heterosis and no indication of a return toward the recurrent parent. The variances were somewhat erratic, but, in general, they were reduced with additional backcrossing. Height and weight-per-volume variances were reduced very little until the Bc_4 , while date of heading and yield variances were higher in the Bc_3 than in the Bc_2 or Bc_4 .

Selection for lines with heading dates like the recurrent parent would have been effective in causing all quantitative characters, except weight-per-volume, to return to the recurrent parent phenotype more rapidly than if random crosses had been made. There was no apparent beneficial or detrimental linkage drag associated with the rust resistance gene contributed by the non-recurrent parent.

The means and variances were used to indicate the type of gene action involved in the determination of each character. Date of heading and yield appeared to result from epistatic gene action, whereas weight-per-volume and height resulted from dominance and additive gene action.

LITERATURE CITED

1. Briggs, F. N. Breeding wheats resistant to bunt by the backcross method. Jour. Amer. Soc. Agron. 22: 239-244. 1930.
2. _____. The backcross method of plant breeding. Jour. Amer. Soc. Agron. 27: 971-973. 1935.
3. _____. The use of the backcross in crop improvement. Amer. Nat. 72: 285-292. 1938.
4. _____ and Allard, R. W. The current status of the backcross method of plant breeding. Agron. Jour. 45: 131-138. 1953.
5. Burnham, C. R. and Cartledge, J. L. Linkage relations between smut resistance and semisterility in maize. Jour. Amer. Soc. Agron. 31: 924-933. 1939.
6. Emsweller, S. L. and Jones, H. A. The inheritance of resistance to rust in the snapdragon. Hilgardia 8: 197-210. 1934.
7. Everson, E. H. and Schaller, C. W. The genetics of yield differences associated with awn barbing in the barley hybrid (Lion x Atlas¹⁰) x Atlas. Agron. Jour. 47: 276-279. 1955.
8. Frey, K. J. Yield components in oats III. Their contribution to the variety x location interaction for grain yield. [To be published in Agron. Jour. ca. 1960].
9. Harlan, H. V. and Pope, M. N. The use and value of backcrosses in small grain breeding. Jour. of Heredity 13: 319-322. 1922.
10. Harrington, J. B. Cereal breeding procedures. FAO [Food and Agriculture Organization] Development Paper No. 28: 16-19. 1952.
11. Hayes, H. K., Immer, F. H., and Smith, D. C. Method of plant breeding. 2nd ed. New York, N. Y., McGraw-Hill Book Co., Inc. 1955.
12. _____, Rinke, E. H., and Tsiang, Y. S. Experimental study of convergent improvement and backcrossing in corn. Minn. Agr. Expt. Sta. Tech. Bul. 172. 1946.
13. Jennings, H. S. The numerical results of diverse systems of breeding. Genetics 1: 53-89. 1916.
14. Johnson, I. J. and Hayes, H. K. The inheritance of pericarp tenderness in sweet corn. Jour. Amer. Soc. Agron. 30: 220-231. 1938.

15. Knight, R. L. Theory and application of the backcross technique in cotton breeding. *Jour. Genetics* 47: 76-86. 1945.
16. Mather, K. The measurement of linkage in heredity. London, Methuen and Co. 1938.
17. _____. Biometrical Genetics. New York, N. Y., Dover Publications, Inc. [c1949.]
18. Mendel, Gregor. Experiments in plant-hybridisation (Translated title). In Sinnott, Edmund W., Dunn, L. C., and Dobzhansky, Th. Principles of genetics. 4th ed. pp. 463-483. New York, N. Y., McGraw-Hill Book Co., Inc. 1950.
19. Murphy, R. P. Convergent improvement with four inbred lines of corn, *Jour. Amer. Soc. Agron.* 34: 138-150. 1952.
20. Nishiyama, I. Cytogenetical studies in *Avena*. I. Chromosome association in hybrids between *A. barbata* Pott. and autotetraploids of *A. strigosa* Schreb. *Cytologia* 7: 276-281. 1936.
21. Powers, Leroy. Studies on the nature of the interactions of the genes differentiating quantitative characters in a cross between *Lycopersicon esculentum* and *L. pimpinellifolium*. *Jour. Genetics* 39: 139-170. 1939.
22. _____. Inheritance of quantitative characters in crosses involving two species of *Lycopersicon*. *Jour. Agr. Res.* 63: 149-174. 1941.
23. Richey, F. D. The convergent improvement of selfed lines of corn. *Amer. Nat.* 61: 430-449. 1927.
24. _____ and Sprague, G. F. Experiments on hybrid vigor and convergent improvement in corn. U.S. Dept. of Agr. Tech. Bul. No. 267. 1931.
25. Riddle, O. C. and Baker, G. A. Biases encountered in large-scale yield trials. *Hilgardia* 16: 1-14. 1944.
26. Ross, W. M. and Miller, John D. A comparison of hill and conventional yield tests using oats and spring barley. *Agron. Jour.* 47: 253-255. 1955.
27. Sampson, D. R. On the origin of oats. Harvard Univ. Bot. Museum Leaflets No. 16: 265-303. 1954.
28. Simpson, D. M. and Weindling, Richard. Bacterial blight resistance in a strain of stoneville cotton. *Jour. Amer. Soc. Agron.* 38: 630-635. 1946.

29. Smith, Harold H. Differential photoperiod response resulting from interspecific gene transfer. *Amer. Jour. Bot.* 36: 828-829. 1949.
30. Suneson, Coit A. An evaluation of nine backcross derived wheats. *Hilgardia* 17: 501-510. 1947.
31. ———, Riddle, O. C., and Briggs, F. N. Yields of varieties of wheat derived by backcrossing. *Jour. Amer. Soc. Agron.* 33: 835-840. 1941.
32. ———, Schaller, C. W., and Everson, E. H. An association affecting yield in barley. *Agron. Jour.* 44: 584-586. 1952.
33. Thomas, Mary. Backcrossing. Cambridge, England. Commonwealth [of Great Britian] Bureau of Plant Breeding and Genetics. 1952.
34. Vallean, W. D. Breeding tobacco varieties resistant to mosaic. *Phytopathology* 36: 412. 1946.
35. ———. Combining resistance to wildfire, mosaic, black root rot, and Fusarium wilt in burley tobacco. *Phytopathology* 38: 27. 1948.
36. Ware, J. O. Plant breeding and the cotton industry. U.S. Dept. of Agr. Yearbook of Agriculture 1936: 657-744. [1936]
37. Wright, Sewell. Systems of mating II. The effects of inbreeding on the genetic composition of a population. *Genetics* 6: 124-243. 1921.

ACKNOWLEDGMENT

The author wishes to express his sincere appreciation to Dr. K. J. Frey for his assistance and guidance during the course of this study and in the preparation of this manuscript.